



User Notes for the PMA Ethiopia Panel Cohort 1 (6-month Follow-up) Survey Dataset, Version 2.0

Disclaimer: PMA cannot provide in-depth support for data analysis or data related questions, however, to assist the end-user, explanation of some variables is provided below.

PMA Ethiopia

Performance Monitoring for Action Ethiopia (PMA Ethiopia) builds on the previous success of PMA2020/Ethiopia and PMA Maternal and Newborn Health study in the Southern Nations, Nationalities and Peoples Region (SNNP). PMA Ethiopia is a five-year project implemented in collaboration with Addis Ababa University, Johns Hopkins University, and the Federal Ministry of Health. It measures key reproductive, maternal and newborn health (RMNH) indicators.

Cross-sectional data, including a health facility-based survey, are collected annually in all regions. Longitudinal data (following pregnant women through one year postpartum) are collected in two cohorts of women (2019-2021 and 2021-2023) in four large, predominantly agrarian regions: Tigray, Oromiya, Amhara, and Southern Nations, Nationalities, and Peoples' Region, and one urban region, Addis Ababa. Afar is included in the first cohort (2019-2021) of the longitudinal survey.

Sampling

PMAET Panel Cohort 1: 6-month Follow-up survey followed women who were eligible to remain in the Panel Cohort 1 study and did not refuse to follow-up at their previous interview. Women who were 5-9 weeks postpartum at baseline who consented to follow-up; women who are pregnant or 0-4 weeks postpartum at baseline, completed the 6-week follow-up survey, and Women pregnant or 0-4 weeks postpartum at baseline, who consented to the 6-week follow up, and had incomplete 6-week forms with results other than "Refused" or "Respondent died" were included in this survey. A total of 2,418 mother or caregivers completed the 6-month follow-up survey. Panel women could become ineligible for 6-month follow-up if they refused follow-up at some point, died, or had no live births.

For more information on the PMA survey methodology and sampling, please refer to PMA Survey Methodology at <u>https://www.pmadata.org/data/survey-methodology</u>.

Materials included in with this data zipped file

The data zipped file includes:

- 1. Panel Cohort 1 6 month follow-up pre-covid questionnaire pdf file
- 2. Panel Cohort 1 6 month follow-up post-covid questionnaire pdf file
- 3. PMAET_Panel_StataSample_Merge_BL_6wkFU_6moFU_v#_Date (Sample Stata dofile)
- 4. Panel Cohort 1 6 month follow-up datasets in 3 formats: csv, xlsx, and dta
- 5. Panel Cohort 1 6 month follow-up datasets in 3 formats: csv, xlsx, and dta
- 6. User notes pdf file

Codebook

The latest version of the PMA Ethiopia Household and Female Survey master codebook can be downloaded from the <u>https://www.pmadata.org/data-codebooks</u>.

Questionnaire Versions

Due to Covid-19 pandemic, the data collection was paused in March 2020 and resumed in late July 2020. When we resumed the data collection, we also fielded a new version of the questionnaire (post-





covid version), so some questions were changed or added. Questions with the suffix _rw were rewritten and questions with the suffix _cc had choice changes.

Six-month Follow-up Survey Specific Information

Timing of Interviews:

We made the decision to interview women outside the 6-7 months postpartum window if their 6month follow-up interview had been delayed due to the pause in data collection. If the timing of interviews matters to your analysis, it's worth paying attention to the elapsed time between the date of interview (**todaySIF** in the dataset) and the delivery date (**recent_birthSIF** in the dataset).

Vaccination dates

For vaccine date analysis, in the 6-month survey we used a different method than baseline and 6-week follow-up to record dates. Instead of having the REs manually enter the date in the Ethiopian calendar, we used an ODK widget that displayed the date in the Ethiopian calendar on the screen but stored the date in Gregorian calendar. This method allowed us to have constraints that kept the dates from being before the birth of the baby or after the interview date.

Additionally, to aid with clarifying the difference between illegible vaccination dates and vaccinations not given, there are a series of questions about the vaccines. VAX stands for each type of vaccine.

VAX_card- Does the vaccine card have VAX and is it legible?

VAX_date_greg- String of the date of VAX for babies where the response to VAX_card was either "Yes, but month or day illegible" or "Yes, legible".

VAX_date_gregSIF- the SIF version of VAX_date_greg. Use this variable for vaccine date analysis

VAX_date_d_dnk- This is a binary variable to indicate if the day in the vaccination date is illegible or unknown. Ex. The RE knew it was in May 2020 but not the day.

VAX_date_m_dnk- This is a binary variable to indicate if the month in the vaccination date is illegible or unknown.

nocard_VAX_yn- If there is no vaccination card, did the baby receive the vaccine?

Dataset Specific Information

Missing Forms:

Some forms did not get submitted or were lost so some of the respondents have a result code of "Missing Form" to indicate that they should have received an interview, but we do not have the form. This particularly affects the Tigray region as data collection was paused in November 2020 due to security concerns.

General Variables

SIF variables: Date and time variables are provided in both string format and as Stata Internal Format (SIF) values. The variable name of any variable that has been changed into SIF is appended with SIF (e.g. **system_date** and **system_dateSIF**). The *"Do not Know"* value for dates is *Jan 1, 2030*. For each





date question where the woman did not know the month but knew the year, the value is Jan 1 of that year and the variable with the same name as the date variable but ending in **_m_dnk** will have a value of 1.

Select multiple variables: Some questions allow for the selection of multiple answers. These variables are in string format and the values are the concatenation of answer choices. Multi-select options are generally, though not always, transformed into binary variables for analysis.

Variable Response Options

Select one: For the select one numeric variables, consistent values for option choices were given across PMA countries. Some select one variables, however, such as **region**, have answer options that vary across countries.

Select multiple: Similarly, most select multiple variables have the same response options across all PMA countries.

See the PMAET HQFQ Master Codebook for complete details on variables and answer choices for each survey.

Specific Variables

EA_ID: The primary sampling unit masked with a random number for anonymity. The same random number is applied to the same EA across multiple survey years.

RE_ID: Identification number of the resident enumerator (RE), or interviewer. RE names are masked with the PMA-Ethiopia random numbers in the household dataset. The same random number is applied to the same REs across different surveys of the PMA-Ethiopia grant.

PMA2020_RE_ID: The resident enumerators (REs), or interviewers, who were involved in the previous PMA2020 surveys (2014-2018), also had PMA2020 IDs. This ID is consistent for all survey rounds of the PMA2020 grant.

QREversion: This variable indicates the version of 6-week follow-up that women received.

SMmetainstanceID: SMmetainstanceID is the unique ID generated by ODK for each 6-month followup form submitted to the central server.

participant_ID: A unique ID of each respondent participating in panel surveys. The same ID is given to the same female respondent across different surveys. This variable should be used to merge or identify women throughout the panel study.

SMFUweight: weight for the 6-month follow-up data.

refuse_future_followup: identifies if the woman refused future follow-up and will not receive the 1-year follow-up

Combining the data with the Baseline Dataset

If you wish to do longitudinal analysis, you will need to merge data from the baseline to the 6-week dataset using **participant_ID**. Example Stata dofiles for merging are included for your reference.





There will be 3 women from the 6-month dataset who do not merge with baseline. Their baseline forms were lost. We include them in cross-sectional analyses of the 6-month data but not longitudinal analyses.

There will be 194 women from baseline who do not merge with the 6-month data. These women became ineligible between screening and baseline, refused follow-up at some point, died, or had no live births.

GPS Variables

GPS coordinates are not released in this dataset.

Notes for Missing Data

In Stata, Missing data is expressed as "." in the cell. Generally, Stata commands perform computations of any type handle missing data by omitting the row with the missing values. However, this may vary across commands. PMA does not impute missing values. Missing data in datasets should be studied and/or treated before proceeding to analysis.

Reasons for missing data:

Normal situations:

- 1. Incomplete forms: If a household, female, or SDP form is not marked as completed (HHQ_result, FRS_result, SDP_result not equal to 1), the observation is likely to miss most of the information. Incomplete forms should not be included in the analysis.
- 2. Observations that are ineligible for subsequent forms: Only eligible respondents will receive subsequent forms. For example, males and ineligible females will not receive female questionnaires in family planning surveys, hence their observations will have all missing values in female forms.
- 3. Question not administered due to skip logic: PMA surveys use ODK's skip logic function. The subsequent questions are administered selectively based on the respondent's previous answers. Irrelevant or inapplicable questions are skipped. For example, a woman who is not a contraceptive user will not be asked questions about contraceptive usage subsequently.

Uncommon situations:

- 1. Lost forms: Due to technical constraints in some challenging data collection areas, forms can be lost in the process of data submission. Although most forms were recoverable, there are occasionally a few that cannot be found. For example, an observation from an eligible woman with completed female form information but missing household form information, or vice versa. These observations may be dropped based on analysis needs.
- 2. Missing due to incorrect skip logic: PMA surveys were conducted under rigorous quality control. However, in rare cases, there can be incorrect skip logic, which skipped a question that was supposed to be administered, resulting in missing values. These errors are documented in the PMA codebook, which can be downloaded from PMA website. It's not necessary to drop the entire observation since this will likely affect only a few questions.



Distinguish missing data from negative values:

- 1. -99: No response. The respondent was administered with the question but did not provide an answer. PMA survey requires consent from the respondent and the respondent has the right to refuse to answer any questions at any point. -99 is recorded to reflect that the respondent did not provide an answer to a certain question.
- 2. -88: Did not know. The respondent consented to answer a specific question but without knowing the answer.
- 3. -77: Not applicable. The question is administered to the respondent but not applicable to the respondent's situation.

Dataset Version Updates

Any updates made to datasets after their initial release will be documented here. PMA releases new versions on as needed basis and users who have been approved previously to get access to the datasets will be notified via email upon the release of the new versions. Users can then log onto the PMA website and re-download the datasets without having to submit a new request.

Users should always be using the latest version of the datasets available on the PMA website. However, if users need an access to the old versions of the datasets for any reasons, users can contact <u>datamanagement@pmadata.org</u> directly with an explanation for why the access to the old version was needed.

In May 2022, a new version of this dataset was released with the following updates:

- Contraceptive calendar variables: calendar_c1_full calendar_c2_full variables were added to the dataset.
- SMFUweight data was updated to be representative of the baseline sample instead of the sample eligible for follow-up. There are no significant changes in estimates using the updated weight compared to the weight from the previous version of the dataset.

Dataset Citations

Suggested citation: Addis Ababa University School of Public Health and The Bill & Melinda Gates Institute for Population and Reproductive Health at The Johns Hopkins Bloomberg School of Public Health. *Performance Monitoring for Action Ethiopia (PMA-ET Panel: Cohort 1 - 6-month Follow-up Survey (Version 2.0), PMAET-Panel-C1-6moFU.* 2021. Ethiopia and Baltimore, Maryland, USA. <u>https://doi.org/10.34976/gvxv-8982</u>

To report errors or inconsistencies:

Please email <u>datamanagment@pmadata.org</u>.